

Recovery  
Oriented  
Language  
Guide

# Contents

- 3 Introduction**
- 4 Recovery perspectives throughout life**
- 5 Guidelines for recovery oriented language**
- 6 Some general guidelines for language and communication
- 8 Out-dated and worn out words
- 11 Key Terms**
- 12 Talking to people at both ends of the age spectrum**
- 12 Young people
- 13 Older people
- 14 A trauma-informed recovery oriented approach**
- 14 Overarching guidelines
- 15 Cultural diversity
- 15 Talking about suicide
- 16 Bibliography and references**



Recovery Oriented Language Guide - Second Edition Revised

© Mental Health Coordinating Council 2018

PO Box 668 Rozelle NSW 2039

**T** 02 9555 8388

**E** [info@mhcc.org.au](mailto:info@mhcc.org.au)

**W** [www.mhcc.org.au](http://www.mhcc.org.au)

***Words are important. The language we use and the stories we tell have great significance to all involved. They carry a sense of hope and possibility or can be associated with a sense of pessimism and low expectations, both of which can influence personal outcomes.***<sup>1</sup>

The Mental Health Coordinating Council (MHCC) developed the ***Recovery Oriented Language Guide*** in 2013 because language matters in all aspects of life. It continues to be particularly important in the context of mental health and recovery. It is vital that words are used that convey hope and optimism, and support and promote a culture that fosters recovery.<sup>2</sup>

People living with mental health conditions are amongst some of the most disadvantaged people in the Australian community, and many live with psychosocial difficulties exacerbated by historical and current trauma, poverty, poor physical health and stigma and discrimination which often feature as part of everyday experiences.<sup>3</sup> The words that we use may effect a person's sense of self and lead to more disadvantage and social exclusion.

The mental health and human service sectors acknowledge MHCC's *Recovery Oriented Language Guide*<sup>4</sup> (the *Guide*) as an extremely useful resource. We have developed this second edition primarily to align with trauma-informed care and practice language approaches, introduce age-related language use and promote the use of supported decision-making language.

Development of the Guide has been informed by a number of sources including: International and Australian literature on trauma-informed and recovery oriented practice; conversations with the mental health practitioners across service sectors; and, most importantly, through listening to the voices of people with lived experience of mental health conditions concerning their recovery journeys.

The Recovery Oriented Language Guide underpins and informs all the work MHCC undertakes in both the policy reform space and in its sector development activities.

You can access this resource from the website at [www.mhcc.org.au/our-work/resources](http://www.mhcc.org.au/our-work/resources)

# Recovery perspectives throughout life

Since 'recovery' was originally defined from the perspectives of adult mental health consumers, MHCC has sought to identify whether these standpoints also apply to people at different developmental stages of life.

The literature suggests that whilst the recovery oriented approach applies to everyone, the language and ways of communicating that approach need to be relevant to particular age groups. MHCC has expanded this edition of the Recovery Oriented Language Guide to include some material concerning the perspectives of young and older people at different stages of their recovery journey. It provides an opportunity for reflection on diversity, which includes young people coming to terms with the new experience of mental health and older people possibly coming to terms with this identity and other associated trauma, grief and loss experiences.

Whatever a person's stage of life, mental health and human services should be familiar with language that reflects a recovery oriented approach to practice, and have an awareness and understanding of the prevalence and impact of trauma, which may have resulted in a range of psychosocial difficulties, and have awareness of the ways in which this may present.

**Cultural diversity considerations** should always guide a worker in their communications.

When a worker is unsure of what is appropriate, they should ask the person what they would like in terms of language interpretation, disability aids, environmental accommodation or supports.

Whatever a person's background, developmental age or impairment experienced, workers should adopt strength-based language to encourage choice and control and support self-directed decision-making.

Just ask a person what they think would assist them achieve their aims and objectives, what they need now and in the future and how they can be supported to communicate and achieve their aims.

Every day we make a countless number of decisions. They are an expression of who we are - our uniqueness, our relationships with others, our achievements and hopes for the future. Sharing power is a key element in promoting recovery.

- Access to supported decision-making can assist a person of any age develop a sense of control over their lives and their recovery. Through decision-making we exercise control over our lives, experience new things and learn about ourselves. Decision-making is so important that it is recognised as a human right
- Decision-making is a skill that can be developed and practised with support.
- Supported decision-making can assist a person to live with meaning, dignity and greater independence.

Appropriate language is a vital component in communicating a sense of self-determination, because feeling powerlessness can be overwhelming, especially when decisions seem to be or are in the hands of others.

Research has shown that communication is only 7 % verbal and 93 % non-verbal. The non-verbal component is made up of body language (55 %) and tone of voice (38 %)⁵ so when communicating with someone it is also important to consider:

Being mindful of the non-verbal aspects of communication, as well as boundaries maintained, appropriate eye contact and using body posture that is non-threatening nor disinterested.

Always try to accommodate a person's developmental age, hearing, cognitive or language difficulties, the time and space to think, question and express their point of view.

It is important to be authentic, transparent and sincere.

**Use plain English language** where possible. Using everyday language can help anyone better understand what is happening to them, their condition, care and treatment or circumstantial change.

**Collaboration and openness** are largely achieved through developing rapport, through connectedness, and a sense of feeling respected and heard.

# Guidelines for Recovery Oriented Language<sup>6</sup>

## General Principles

### The language we use:

- Represents the meanings we have constructed from experience
- Prompts attitudes, expectations and actions
- Should always reflect 'unconditional positive regard'<sup>7</sup> for people.

We may be unaware of how the words we use reflect our attitudes and the impact they have upon those around us.

The words we choose reflect our attitudes - that we do (or do not) truly value people, believe in and genuinely respect them.

None of us should be defined by the mental health conditions or psychosocial difficulties that we experience, or by any single aspect of who we are; we should be respected as individuals first and foremost.

### Our language needs to be:

- Respectful
- Non-judgemental
- Clear and understandable
- Free of jargon, confusing data, and speculative comment
- Consistent with our body language
- Sincere in carrying a sense of commitment, hope and presenting the potential for opportunity
- Strengths-based

### We need to give thought to:

- How the language we use, is read and heard by the person to which we are communicating, and how it may positively or negatively contribute to their health and wellbeing
- What meanings we present to people to live by

Our language conveys our thoughts, feelings, facts and information, but beyond that, we need to be reflective in our practice and ask ourselves questions like:

- *What else am I saying?*
- *How will someone else read or hear this?*
- *Do I give a sense of commitment, hope and present opportunity or a sense of pessimism?*
- *Do I convey an awareness and expectation of recovery?*

The approach to language when talking to people needs to take into consideration where they are in their recovery journey. This may fluctuate in relation to their physical and mental health, and social and emotional wellbeing.

## Some general guidelines for language and communications<sup>8</sup>

### DO

**DO** put people first:

**DO** say “person with mental health condition”

**DO** say “a person who has been diagnosed with...”

**DO** emphasise abilities

*Do focus on what is strong  
i.e., the person’s strengths, skills & passions*

**DO** use language that conveys hope and optimism that supports, and promotes a culture of recovery

**DO** enquire as to how the person would like to be addressed

**DO** use language that is comfortable for you and reflects your genuine, true self

**DO** clarify that people understand the information they have been given

Make sure that whatever a person’s age, cultural background and cognitive skills that they have understood what has been said

### DON'T

**DON'T** label people:

**DON'T** say “he is mentally ill”, “she is mentally ill”

**DON'T** define the person by their struggle or distress

**DON'T** equate identity with a person’s diagnosis

*Very often there is no need to mention a diagnosis at all. It is sometimes helpful to use the term “a person diagnosed with”, because it shifts the responsibility for the diagnosis to the person making it, leaving the individual the freedom to accept it or not.*

**DON'T** emphasise limitations.

*Don’t focus on what is (in your mind) wrong with the person*

**DON'T** use condescending, patronising, tokenistic, intimidating or discriminating language

**DON'T** make assumptions based on external appearances or communication difficulties

**DON'T** sensationalise a mental illness  
*This means not using terms such as “afflicted with,” “suffers from” or “is a victim of”*

**DON'T** portray successful persons with mental health conditions as superhuman. *This carries the assumption that it is rare for people with a mental health condition to achieve great things*

**DON'T** presume that a person wants to be called by a particular term (e.g., consumer or client) and check whether the wish to be addressed by their family or first name (e.g., Ms Smith or Kylie) or another name which they identify

**DON'T** use jargon, or unfamiliar language.

**DON'T** use specialist or medical language unless you accompany it with plain English explanations

DO	DON'T
DO use language that conveys optimism and positivity	DON'T use negative or judgemental language
DO ask <i>“what is important to you?”</i>	DON'T refer negatively to aspirations identified in the past that a person did not follow up
DO ask <i>“what are you looking forward to doing?”</i>	
DO ask <i>“what do you think might be steps forward”</i>	DON'T use the concept of goals with young people or older people unless it feels appropriate. Rather talk about aspirations, dreams and hopes
DO ask whether the person feels they have been consulted and listened to about their care, treatment or support plans	DON'T argue with a person's perception of events
DO validate a person's experiences	DON'T minimise a person's experience in the urgency of managing symptoms
DO ask whether the person has been given the opportunity to ask questions, and check that they have the information they need	DON'T argue that information was already provided or known
DO check that an older person has heard and understood what has been said clearly - when you know or sense they may have hearing and/or cognitive difficulties	DON'T assume that having said something, that it is understood
DO allow people the time to find the words and express what they need to say	DON'T jump in and speak for someone
DO ask people if they feel ready to make their own decisions or would like to be supported, and in what way	DON'T tell someone that certain information is irrelevant
DO ask what has been helpful and unhelpful in the past	DON'T harp on failures of the past
DO involve people in the development of treatment, care and support planning	DON'T assume that you know what is best for a person
DO involve others providing care coordination across services	DON'T devise treatment, care or a support plan without consultation with the person you are working with
DO be mindful of the importance of individual identity to all people, but be particularly sensitive to peoples' fears of being considered to lack decision-making capacity	DON'T make assumptions about people based on their diagnosis
DO be mindful of older people's fear of losing their sense of identity or a young person being considered mature enough to make decisions	DON'T make assumptions about age or disability. Remember older people have a lifetime of experience, and many young people have roles of responsibility despite their age

## Out-dated and worn-out words<sup>9</sup>

Language of Acceptance, Hope, Respect & Uniqueness	Worn-out words
<ul style="list-style-type: none"> <li>■ Kylie has a mental health condition or a disability</li> </ul>	<ul style="list-style-type: none"> <li>■ Kylie is not normal</li> </ul>
<ul style="list-style-type: none"> <li>■ Sam lives with a mental health condition</li> <li>■ Sam is a person with lived experience of ....</li> <li>■ Sam has been told he has schizophrenia</li> <li>■ Sam has been diagnosed with .....</li> <li>■ Sam has experienced anorexia</li> <li>■ Sam is a person with lived experience of trauma</li> <li>■ Sam is a person who experiences mental health and co-existing difficulties with substance use</li> </ul>	<ul style="list-style-type: none"> <li>■ Sam is mentally ill</li> <li>■ Sam is schizophrenic</li> <li>■ Sam is a bipolar</li> <li>■ Sam is an anorexic</li> <li>■ Sam has PTSD</li> <li>■ Sam is a PD</li> <li>■ Sam is a borderline</li> <li>■ Sam has drug problems</li> </ul>
<ul style="list-style-type: none"> <li>■ Kylie is having a rough time</li> <li>■ Kylie is having difficulty with her recommended medication</li> <li>■ Kylie's medication is not helping her</li> <li>■ Kylie is experiencing unwanted effects of her medication</li> <li>■ Kylie disagrees with her diagnosis</li> <li>■ Kylie is experiencing ...</li> </ul>	<ul style="list-style-type: none"> <li>■ Kylie is decompensating</li> <li>■ Kylie is treatment resistant</li> <li>■ Kylie is uncooperative</li> <li>■ Kylie doesn't accept she is mentally ill</li> <li>■ Kylie has no insight</li> </ul>
<ul style="list-style-type: none"> <li>■ Sam is trying really hard to self-advocate and get his needs met</li> <li>■ Sam may need to work on more effective ways of getting his needs met</li> </ul>	<ul style="list-style-type: none"> <li>■ Sam is manipulative, irritable</li> <li>■ Sam is demanding and unreasonable</li> <li>■ Sam has challenging or complex behaviours</li> <li>■ Sam is dependent</li> </ul>
<ul style="list-style-type: none"> <li>■ Kylie is choosing not to...</li> <li>■ Kylie would rather look for other options</li> </ul>	<ul style="list-style-type: none"> <li>■ Kylie is non-compliant</li> <li>■ Kylie has a history of non-compliance</li> </ul>
<ul style="list-style-type: none"> <li>■ Sam is pleased or satisfied with the plan we've developed together</li> <li>■ Sam and the team have developed a good rapport</li> <li>■ Sam is able to seek help and recognises when things are not going so well</li> <li>■ Sam is working hard towards achieving his goals</li> <li>■ Sam is taking each day at a time</li> </ul>	<ul style="list-style-type: none"> <li>■ Sam is compliant or manageable</li> <li>■ Sam has partial insight</li> <li>■ Sam is cooperating</li> <li>■ Sam has acquired insight</li> <li>■ Sam is learning to manage his illness</li> <li>■ Sam is unmotivated</li> </ul>



## Language of Acceptance, Hope, Respect & Uniqueness

## Worn-out words

- Kylie chooses not to...
- Kylie is concerned about the health implications of her treatment plan
- Kylie prefers not to...
- Kylie is very independent
- Kylie seems unsure about...
- Kylie might benefit from some help at home

- Kylie has no insight
- Kylie is treatment resistant
- Kylie refuses support
- Kylie won't engage with services
- Kylie needs support with her ADLs
- Kylie is low functioning

- Sam is really good at...

- Sam is high functioning

- Kylie has a tough time taking care of herself
- Kylie has a tough time learning new things
- Kylie is still considering her options
- Kylie is still working out what she needs

- Kylie is low functioning

- Sam tends to (describe actions, e.g., hit people) when he is upset
- Sam sometimes kicks people when he is hearing voices
- Sam is finding it difficult to socialise
- Sam likes his own company

- Sam is dangerous; abusive; angry, aggressive
- Sam demonstrates challenging, high risk behaviour/s
- Sam is high risk
- Sam is anti-social
- Sam is isolative
- Sam doesn't want to socialise

- Kylie is experiencing both mental health and substance use problems
- Kylie tends to use non-prescribed substances to help manage distress/cope with life

- Kylie is dually diagnosed
- Kylie has comorbidities or is comorbid
- Kylie is MICA/MISA (mentally ill chemically abusing, mentally ill substance abusing)
- Kylie is an addict

- Sam doesn't seem ready to go back to work
- Sam is not in an environment that motivates him
- Sam is working on finding his motivation
- Sam has not yet found anything that sparks his interests

- Sam is unmotivated
- Sam is not engaged or does not want to be engaged
- Sam isolates
- Sam rejects help

- Kylie has a lot of energy right now
- Kylie hasn't slept in three days
- Sam is experiencing a lot of fear
- Sam is worried that his neighbours want to hurt him
- Sam often disagrees and gets angry with his family, friends etc
- Kylie has been working towards recovery for a long time
- Kylie has experienced serious depression for many years
- So far, Kylie has not accessed support that has been helpful to her

- Kylie is manic
- Kylie is hyper
- Sam is paranoid
- Sam is delusional
- Sam is aggressive
- Kylie has a chronic mental illness
- Kylie is severely mentally ill
- Kylie will never recover - she rejects help

- Sam and I aren't quite on the same page
- It is sometimes challenging for me to work with Sam
- Sam has not had good experience with services in the past
- Sam is a young person who has recently been given a diagnosis and is having difficulty to come to terms with this news

- Sam is very difficult
- Sam has challenging behaviour
- Sam won't engage with services
- Sam doesn't accept what he has been told by the treating team

*If worn out words are used to describe people's attempts to reclaim some shred of power while receiving services in a system that may try to control them, then important opportunities to support a person's recovery will be lost.*

*A person trying to get their needs met - may have a perception or opinion different from, or not shared by, others - and their actions may not be effectively bringing them to the results they want.*

- Manipulative
- Grandiose
- In denial
- Passive aggressive
- Self-defeating
- Oppositional
- Personality disordered

- Charles is an older person who displays frustration at times
- Charles is someone used to being independent who is finding it difficult accepting support
- Charles may need some support to help him make decisions
- Charles may need support in some areas of his life
- We need to find out what we can do to support Charles that best suits him

- Charles is a challenging, difficult, grumpy man
- easily angered, irrational and short tempered
- Charles rejects help and advice
- Charles isn't capable of deciding for himself what's best
- Charles has complex needs
- Charles has poor ADLS
- Charles is uncooperative

# Key Terms

**A Consumer** is a person with lived experience of a mental health condition who is accessing or has previously accessed a mental health service.<sup>10</sup> Within a child and youth mental health context, both the parents and the child or young person may sometimes also be described as consumers.

**Capacity** refers to a person's ability to make his/her own decisions. These may be small decisions, such as what to do each day, or bigger decisions such as where to live or whether to have an operation. A person may lack capacity in some areas, but still be able to make other decisions.

**Cognitive functioning** refers to the underlying cognitive processes that allow for effective information processing that assist decision-making, planning and completing actions.

**Complex need** is commonly used to refer to individuals who present with an inter-related mix of diverse mental health and physical health issues, developmental and psychosocial problems.

**Dignity of Risk** refers to the individual's right to make informed choices in relation to a variety of life experiences and take advantage of opportunities for learning, developing competencies and independence and, in doing so, takes a calculated risk.

**Diversity** is inclusive of but not limited to the diversity among people with respect to culture, religion, spirituality, disability, power, status, gender and sexual identity and socioeconomic status.<sup>11</sup>

**Peer Work** is a fast growing occupational group in the mental health workforce. Peer services are a core component of a genuinely recovery based service. Peer work, peer workers and peer workforce includes all workers in mainstream or alternative mental health services or initiatives who are employed to openly identify and use their lived experience of mental illness and recovery as part of their work. Peer support workers provide support for personal and social recovery to other people with mental health conditions, including in acute mental health settings housing, supported employment, community-based support and so on.<sup>12</sup>

**Psychosocial disability** is an internationally recognised term under the United Nations Convention on the Rights of Persons with Disabilities. It is characteristically used to describe many negative outcomes for a person living with a mental health condition attempting to interact with a social environment that presents barriers to their equality with others. Psychosocial disability may also describe the experience of people regarding participation restrictions related to their mental health conditions as the loss of or reduced abilities to function, think clearly, experience full physical health and manage the social and emotional aspects of their lives.

**Recovery** is defined as being able to create and live a meaningful and contributing life in a community of choice with or without the presence of mental health issues.<sup>13</sup>

**Somatisation** is a tendency to experience and communicate psychological distress in the form of somatic symptoms and behavioural features. More commonly expressed, it is the generation of physical symptoms of a psychiatric condition such as anxiety.

**Supported decision-making** is an approach designed to support people make significant decisions, exercise their legal capacity, make day-to-day choices and draw upon their strengths and support networks.

**Trauma-informed** is the re-conceptualisation of traditional approaches to health and human service delivery whereby all aspects of services are organised acknowledging the prevalence of trauma throughout society. 'Trauma-informed' services are aware of and sensitive to the dynamics of trauma as distinct from directly treating trauma per se.

# Talking to people at both ends of the age spectrum

## Young People

Just like with people from other groups or diverse cultures, when communicating with a young or older person using language that they are comfortable with is important. Be guided by their style of speech.

Young people are no different to adults in expressing the importance of collaboration and openness as worker attributes. What is particularly meaningful is displaying a genuine interest in people and their lives. Asking “how’s everything going?” can be a good way of opening the door to a conversation about anything that they may need or like to talk about. Young people characteristically relate more to concepts of health and wellbeing rather than illness and recovery.

**Use language that is real and familiar rather than imitate young people.**

The idea of being asked to formulate recovery goals, particularly for young people can lead them to feel judged, especially when they are unable to list concrete objectives.

Instilling hope is vital to everyone, but some young people feel overwhelmed thinking of the future. Talk instead about hopes and dreams that may have for themselves.

*Workers should not feel uncomfortable or needing to use language that they wouldn't normally use. However it is important to understand that swearing and 'bad language' is a prominent feature in the vocabulary of many young people - both when things are going well and when things are challenging.*

A young person may say “I feel crap” or “this is shit”. Enabling conversation that is accepting of this language is important in establishing rapport with a young person. Some of the expressions used by young people may offend others from different age groups and cultures, nevertheless, it important to be accepting of contemporary vernacular.

*An alternative approach is to refer to 'steps forward' rather than 'goals'. For example, "what do you think may be some useful steps forward?" or "what are you looking forward to doing (e.g. when you are discharged from hospital, go home etc.)?" Young people are often figuring out who they are and what they want of life and don't want to be cornered.*

Young people are increasingly comfortable using technology-based communications to discuss their emotions and experiences.<sup>14</sup> It would appear that the absence of social cues such as facial expressions and gestures provides young people with an opportunity to disclose serious or sensitive information in what they perceive as a less-judgmental environment (where they can meet and converse with like-minded people).<sup>15</sup> Health practitioners have noted that meeting young people in a space where they are comfortable can help build rapport and improve communication, even when online.<sup>16</sup> Having discarded the formalities of meeting face-to-face, online communication can offer a vehicle for frank and sincere discussion about a person’s mental health difficulties.

***The NSW Child and Adolescent Mental Health Services Competency Framework<sup>17</sup> identifies the importance of mental health workers being culturally sensitive to adolescents when working with them; for example, appropriate non-verbal communication, eye contact and body posture.***

## Older persons

**Using a recovery oriented approach** sets out to enable a person living with a mental health condition to create and live a meaningful and contributing life in their community of choice.<sup>18</sup>

Many older people have a clear sense of who they are and how they define themselves, and can build on a lifetime's experience and resilience which can buffer the impact of any illness or circumstance.

**Working with older people often includes:**

**Supporting** a person to maintain a sense of enduring self-identity. Older people living with mental health conditions have described this as 'continuing to be me'.<sup>19</sup> It is important to express a genuine interest in knowing who they are.

**Understanding** that for some older people's mental health conditions may have become so entrenched that their sense of who they are is compromised by illness and its impacts. This may result in them appearing helpless and hopeless. The language we use should reflect the fact that there are other perspectives without minimising what they are actually feeling.

**Awareness** that older people tend to conceptualise recovery rather differently to young people and adults more generally. Their aim may be simply to maintain a sense of who they are despite the disabilities they experience, and be valued and respected to know what they want and need.

**Considering** that a person may feel patronised and cornered when asked about goals for recovery. Older people may prefer to have their recovery journey expressed in a way which focuses on what will give them the best life they can live in the circumstances.

There is evidence suggesting that older people's dignity and autonomy is being undermined in health care settings and that a sizable cross section of healthcare professionals hold stereotypical, negative attitudes towards older people.<sup>20</sup> Diminishing mental health among older people is often not identified by relatives, health care professionals and older people themselves who may attribute symptoms of to the effects of ageing or to physical and environmental changes.

Often service providers make assumptions about older people, and what is appropriate communication. Important to language in this context is communicating respect and supporting choice and autonomy whatever difficulties a person may be having. Importantly, assuming capacity as a first principle.<sup>21</sup>

***Three components of recovery appear to be distinct to older people: the significance of an established and enduring sense of identity; coping strategies which provide continuity and compensation and therefore reinforce identity, and the impact of coexisting physical illness.***

***It is vital that when communicating with an older person that they are supported to have their voice heard and their choices understood. Older people should be reassured that their autonomy and ability to self-determine life choices will not be undermined unnecessarily, especially when other disabilities may be involved.***

***Avoid asking others, even those close to them about what they want, unless a person clearly wants someone else to speak on their behalf, or are unable to communicate their preferences.***

# A trauma-informed recovery-oriented approach

Recovery orientation has been adopted as an overarching philosophy to guide mental health practice and is embedded into policy and standards nationally.<sup>22</sup> An understanding of trauma is integral to a recovery-oriented approach. In fact developing and implementing trauma-informed systems of care is one of the first steps towards becoming recovery oriented. Critical to this objective is to use language that reduces the possibilities for re-traumatisation and harm within service systems and practice. A trauma-informed recovery-oriented approach is person centred and involves sensitivity to individuals' particular needs, preferences, safety, vulnerabilities and wellbeing, recognises lived experience and empowers people with lived experience to genuinely participate in decision-making.

## Overarching guidelines

- Speak or write about a person with a mental health condition, psychosocial disability, cognitive impairment, problem and/or difficulty; not about a disorder, diagnosis, symptom/s and/or case or bed or a derisory term such as 'frequent flyer' or 'blocked bed'.
- Always include a description of a person's strengths and resourcefulness alongside the difficulties they experience.
- Where applicable, explicitly own words and concepts such as diagnosis or assessment as from a medical/service provider opinion or perspective rather than as a pronouncement of universal truth.
- Do not make assumptions and describe achievements, possessions or connections as merely grandiose delusions.
- Do not assume that disclosures of abuse are necessarily imaginary or represent part of the psychosis a person may be experiencing.
- Do not assume that risks presented in files and notes that relate to the distant or very distant past have current relevance.
- Record people's progress and their efforts towards their own recovery, the steps forward that they have made, using the person's own words and meaning.
- Where there are different views between the person writing a letter or report and the person it is important to:
  - include recognition of that awareness
  - describe their viewpoint in their own words, and
  - describe how their viewpoint contrasts with the author's

For example, "*whereas I think ... I'm aware that Sam has a very different point of view and considers or stated that ...*"

  - Note directions for negotiating these differences
- Express 'shortfalls' as work or progress still to be achieved.
- Record the person's own hopes or ambitions as well as those held by the support or treating team and what needs to happen for such hopes to be realised.
- Seek to express issues of risk (safety appraisal) in terms of planning for recovery, safety and success; including for people who may be required to comply with involuntary treatment.
- Seek to ensure that issues of compromised safety include risk of re-traumatisation as a consequence of a range of involuntary treatment, including detention in a hospital environment.
- When actions are suggested that the person disagrees with, give a clear reason why these are considered necessary in terms of supporting someone's recovery, and acknowledge their alternate view.
- Ensure demonstration of respect with reference to people's concerns about the physical and psychological impacts of medication/s that they are expected to accept, and that discussions appropriately factor in the competing risks.
  - When there is opportunity, such as in regards to Mental Health Review Tribunal determinations, always offer a copy to the person following the hearing decision.
- In respect to reports to be presented to a Tribunal, always offer to discuss the draft prior to a hearing unless there is good reason, in which case:
  - offer to review and respond to their views on what you have written
  - where there are significantly different viewpoints, consider how these can be included either by amending what you have written if it is acceptable to you or otherwise include a description of the person's alternate viewpoint in the file



- Be aware that letters and reports are constructions rather than objective descriptions:
  - where possible, write reports with the person they are about, while at the same time preserving the integrity and authenticity of your own viewpoint
  - Where not possible, write them knowing that the person may read them
- Where there is a practice of offering people copies of letters written about a person, consider if the letter could instead be written directly to the person it is about - as a record of the conversation and a reminder of decisions - and copied to the other relevant parties (e.g., peer workers, support workers, general practitioners).
- Set up recovery-oriented language prompts in organisational documents and data templates, and include in continuous improvement audit processes.
- Ensure that in talking to anyone, that environmental safety has been established before discussing anything that may represent a trauma trigger for a person you are supporting.

## Cultural Diversity

People from different cultures may express their distress in physical or somatic symptoms, or in descriptive terms unusual to you. It is important that workers pay attention to the person's description of their lived experience.

Living with a mental health condition may be considered a weakness in some cultures, and some people may find a diagnosis shaming or guilt provoking, whilst others may consider it a relief to put a name to their difficulties. Try and find out how a person's culture affects the way they perceive their condition and use language that most appropriately relates to their experience.

## Talking About Suicide<sup>23</sup>

One impetus to change the language of suicide began in the bereavement community. In addition to the insensitive language often used to describe suicide, and the silence and denial - the absence of suicide language and conversation is a major contributor to the stigma people face in the community.<sup>24</sup>

Suicide often leaves the bereaved with especially acute feelings of self-recrimination. Those who are left behind may feel the full burden of suicide's stigma, and can feel abandoned and ashamed. Added to this injury is the mention of suicide in euphemistic language that goes to great lengths to neutralise the real meaning that exists concerning death as a consequence of suicide and the loss attached to it. Because this silence can be debilitating, the need for language that addresses the act of suicide in a direct but respectful way was identified and has, in recent years, gathered momentum.<sup>25</sup>

***Suicide is no longer a crime, and so we should stop saying that people commit suicide. We now live in a world where we seek to understand people who experience suicidal thoughts, behaviours and attempts, and then to treat them with compassion rather than condemn them.***<sup>26</sup> Part of this is to use appropriate, non-stigmatising language when referring to suicide.

Often people seek attention from others for comfort and reassurance when they are distressed. In the context of a suicide attempt or suicidal ideation people are often described as 'attention seeking'. This is unhelpful language suggesting that a person is repeatedly displaying negative behaviours to gain attention. It is important to acknowledge the sequence of events leading to the situation arising and then state the actual behaviour of concern.

### Appropriate Words

- Died by suicide
- Took his or her life
- Ended his or her life
- Non-fatal attempt at suicide
- Attempt to end his or her own life

### Worn-out words

- Committed suicide
- Successful suicide
- Completed suicide
- Failed attempt at suicide
- Unsuccessful suicide

## Bibliography

Alberta Mental Board 2005, 'What's in a word: the language of suicide', Available: <http://www.albertahealthservices.ca/MentalHealthWellness/hi-mhw-sps-language-of-suicide-1p.pdf>

Beaton, S Forster P & Maple M 2013, 'Suicide and language: Why we shouldn't use the 'C' word', *InPsych*, v. 35 (1), p. 30-31 (2013) -- issn: 1441-8754 -- Australian Psychological Society.

Brooker D 2008, 'Person centred care', In: Jacoby R, Oppenheimer C, Dening T & Thomas A, (eds) Oxford textbook of old age psychiatry, Oxford: Oxford University Press.

beyondblue 2018, 'Language when talking about suicide', Available: <https://www.beyondblue.org.au/the-facts/suicide-prevention/worried-about-suicide/having-a-conversation-with-someone-you%27re-worried-about/language-when-talking-about-suicide>

Commonwealth of Australia 2013, 'National framework for recovery-oriented mental health services'.

Daley, S, Newton, D, Slade, M, Murray, J & Banerjee, S 2012, 'Development of a framework for recovery in older people with mental disorder', *International Journal of Geriatric Psychiatry*, 28 pp. 522-529

Department of Health and Ageing 2012, 'National Recovery Oriented Mental Health Practice Framework', Commonwealth of Australia

Devon Partnership Trust and Torbay Care Trust 2008, 'Putting Recovery at the Heart of All We Do', UK.

Lothian, K & Philp, I 2001, 'Maintaining the dignity and autonomy of older people in the healthcare setting', *BMJ* 322. 668-670.

Mental Health Drugs and Regions Division 2012, 'Framework for recovery-oriented practice: Literature review', Victorian Government Department of Health, Melbourne.

Mental Health Coordinating Council 2013, 'Trauma-Informed Care and Practice: Towards a cultural shift in policy reform across mental health and human services in Australia, A National Strategic Direction', Position Paper and Recommendations of the National Trauma-Informed Care and Practice Advisory Working Group, Authors: Bateman, J & Henderson, C (MHCC) Kezelman, C (Adults Surviving Child Abuse, ASCA). Available: [http://www.mhcc.org.au/media/32045/ticp\\_awg\\_position\\_paper\\_\\_v\\_44\\_final\\_\\_07\\_11\\_13.pdf](http://www.mhcc.org.au/media/32045/ticp_awg_position_paper__v_44_final__07_11_13.pdf)

Mental Health Coordinating Council 2013, 'Recovery Oriented Service Self-Assessment Toolkit' (ROSSAT) Implementation Project: Literature Review.

Mental Health Coordinating Council 2014, 'Recovery Orientation in Youth Mental Health and Child and Adolescent Mental Health Services (CAMHS)', NSW, Australia. Available: <http://www.mhcc.org.au/media/50501/mhccrecoveryforyoungpeople-discussionpaper.pdf>

Mental Health Coordinating Council 2015, 'Youth Recovery Language Project, Literature Scan and Report'.

NSW Consumer Advisory Group - Mental Health Inc. and Mental Health Coordinating Council 2014, 'Recovery Oriented Service Self-Assessment Toolkit (ROSSAT): A Recovery Resource for Mental Health Community Managed Organisations Project - Final Project Report'.

NSW Health Illawarra Shoalhaven LHD 2016, 'Clinical Support Tool: Guide to Appropriate Language in Mental Health Services', Available: [http://www.mhrt.qld.gov.au/wp-content/uploads/2011/10/guide-to-Appropriate-Language-policy\\_1.pdf](http://www.mhrt.qld.gov.au/wp-content/uploads/2011/10/guide-to-Appropriate-Language-policy_1.pdf)

NSW Ministry for Health 2011, 'NSW Child and Adolescent Mental Health Services (CAMHS) Competency Framework, Competency 5'. Available: <http://www.health.nsw.gov.au/mentalhealth/programs/mh/Publications/camhs-nov11.pdf>

National Mental Health Consumer & Carer Forum 2011, 'Unravelling Psychosocial Disability: A Position Statement by the National Mental Health Consumer & Carer Forum on Psychosocial Disability Associated with Mental Health Conditions'.

Roberts, G & Thekkepalakkal, A 2009, 'Developing Recovery Oriented Practice - A guide to writing reports and letters: Recovery and Independent Living', PEG Advisory Paper 9, Devon Partnership Trust, UK.

Rogers, C 1951, "Client-centered Approach to Therapy", in I. L. Kutash and A. Wolf (eds.), *Psychotherapist's Casebook: Theory and Technique in Practice*. San Francisco: Jossey-Bass.

Slade, M 2009, '100 Ways to Support Recovery: A Guide for Mental Health Professionals'. UK

Wahl O 1999, 'Mental health consumers' experiences with stigma', *Schizophrenia Bulletin*, 25, 467-478, USA.

## References

1. Devon Partnership Trust and Torbay Care Trust 2008, 'Putting Recovery at the Heart of All We Do', UK, p. 2.
2. Department of Health & Ageing 2012, 'National Recovery Oriented Mental Health Practice Framework'.
3. National Mental Health Consumer & Carer Forum (NMHCCF) 2011 'Unravelling Psychosocial Disability: A Position Statement by the National Mental Health Consumer & Carer Forum on Psychosocial Disability Associated with Mental Health Conditions'.
4. MHCC 2013, Recovery Oriented Language Guide, First edition © Mental Health Coordinating Council 2013.
5. Mehrabian, A 1972, *Nonverbal Communication*. New Brunswick: Aldine Transaction.
6. Adapted from Roberts, G & Thekkepalakkal, A 2009, *Developing Recovery Oriented Practice - A guide to writing reports and letters: Recovery and Independent Living PEG Advisory Paper 9*. Devon Partnership Trust, UK.
7. Rogers, Carl 1951, "Client-centered Approach to Therapy", in I. L. Kutash and A. Wolf (eds.), *Psychotherapist's Casebook: Theory and Technique in Practice*. San Francisco: Jossey-Bass.
8. Adapted from: Wahl O 1999, 'Mental health consumers' experiences with stigma', *Schizophrenia Bulletin*, 25, 467-478, USA.
9. Ibid.
10. Australian Health Ministers 2003, 'National Mental Health Plan 2003-2008', Commonwealth of Australia, Canberra, ACT.
11. State of Victoria, Department of Health 2013, 'National Practice Standards for the Mental Health Workforce 2013', Victorian Government Department of Health, Melbourne, VIC.
12. Interrelate 2013, 'Peer Workforce in Mental Health: Part 1: Proposal to Develop an International Consensus to the International Initiative for Mental Health Leadership (IIMHL)', National Empowerment Center (NEC), Lawrence, MA.
13. Commonwealth of Australia 2013, 'A national framework for recovery-oriented mental health services: Policy and theory', Australia, p. 17.
14. Eysenbach G Powell J Englesakis M Rizo C & Stern A 2004, 'Health related virtual communities and electronic support groups: systematic review of the effects of online peer to peer interactions', *British Medical Journal*, vol. 328, no. 7449, pp. 1166.
15. Campbell AJ & Robards F 2012, 'Using technologies safely and effectively to promote young people's wellbeing: a better practice guide for services', NSW Centre for the Advancement of Adolescent Health, Westmead and the Young and Well Cooperative Research Centre, Abbotsford.
16. Ibid.
17. NSW Ministry for Health, 2011, 'NSW Child and Adolescent Mental Health Services (CAMHS) Competency Framework, Competency 5', p. 5.
18. Commonwealth of Australia 2013, 'National framework for recovery-oriented mental health services'.
19. Daley, S, Newton, D, Slade, M, Murray, J & Banerjee, S 2012, 'Development of a framework for recovery in older people with mental disorder', *International Journal of Geriatric Psychiatry*, 28 pp. 522-529
20. Lothian, K & Philp, I 2001, 'Maintaining the dignity and autonomy of older people in the healthcare setting', *BMJ* 322. 668-670.
21. Brooker D 2008, 'Person centred care', In: Jacoby R, Oppenheimer C, Dening T & Thomas A, (eds) Oxford textbook of old age psychiatry, Oxford: Oxford University Press.
22. Australian Health Ministers' Advisory Council, 2013, A national framework for recovery-oriented mental health services: Guide for practitioners and providers, Commonwealth of Australia. Available: [https://www.health.gov.au/internet/main/publishing.nsf/content/67D17065514CF8E8CA257C1D00017A90/\\$File/recovgde.pdf](https://www.health.gov.au/internet/main/publishing.nsf/content/67D17065514CF8E8CA257C1D00017A90/$File/recovgde.pdf) [Accessed March 2018]
23. Beaton, S Forster P & Maple M 2013, 'Suicide and language: Why we shouldn't use the 'C' word', *InPsych*, v. 35 (1), p. 30-31 (2013) issn: 1441-8754, Australian Psychological Society.
24. beyondblue, 2018, Quote: Susan Beaton, Suicide prevention specialist, 'Language when talking about suicide', Available: <https://www.beyondblue.org.au/the-facts/suicide-prevention/worried-about-suicide/having-a-conversation-with-someone-you%27re-worried-about/language-when-talking-about-suicide> [ Accessed March 2018]
25. Alberta Mental Board, 2005, 'What's in a word: the language of suicide', [Accessed July 2017] Available:
26. <http://www.albertahealthservices.ca/MentalHealthWellness/hi-mhw-sps-language-of-suicide-1p.pdf>
27. beyondblue, 2018, Quote: Susan Beaton, Suicide prevention specialist, 'Language when talking about suicide', Available: <https://www.beyondblue.org.au/the-facts/suicide-prevention/worried-about-suicide/having-a-conversation-with-someone-you%27re-worried-about/language-when-talking-about-suicide> [ Accessed March 2018]